

Intractability Rate of Approximation Problem for Random Fields in Increasing Dimension

Nora Serdyukova

December 11, 2006

Abstract

The behavior of average approximation cardinality for d -parametric random fields of tensor product type is investigated. It was shown in [5] that for given relative error level the approximation cardinality increases exponentially, as $d \rightarrow \infty$, i.e. one observes the curse of dimensionality (intractability) phenomenon.

The aim of this paper is to give the exact asymptotic expression for the approximation cardinality.

Key words: tractability, random fields, Gaussian processes, linear approximation error, complexity, curse of dimensionality.

1 Introduction

Suppose we have a random function $X(t)$, with t in some parametric set T , admitting a series representation via random variables ξ_k and the deterministic real functions φ_k , namely

$$X(t) = \sum_{k=1}^{\infty} \xi_k \varphi_k(t).$$

For any finite set of positive integers $K \subset \mathbb{N}$ let $X_K(t) = \sum_{k \in K} \xi_k \varphi_k(t)$. In many problems one needs to approximate X with finite rank process X_K . Natural questions arise then: how large should be K that yields a given small approximation error? Given the size of K , which K provides the smallest error? In this article, we address the first of these questions for a specific class of random functions, namely, *tensor product type random fields* with high-dimensional parameter sets.

Let $(\lambda(i))_{i \geq 1}$ be a non-negative sequence satisfying

$$(1.1) \quad \sum_{i=1}^{\infty} \lambda(i)^2 < \infty$$

and let $(\varphi_i)_{i>0}$ be an orthonormal system in $L_2[0, 1]$.

Consider a family of tensor product type random fields $\mathbb{X} = \{X^{(d)}(t), t \in [0, 1]^d\}$, $d = 1, 2, \dots$, given by

$$(1.2) \quad X^{(d)}(t) = \sum_{k \in \mathbb{N}^d} \xi_k \prod_{l=1}^d \lambda(k_l) \prod_{l=1}^d \varphi_{k_l}(t_l),$$

where ξ_k are non-correlated random variables with zero mean and unit variance. Obviously, under assumption (1.1) the sample paths of $X^{(d)}$ belong to $L_2([0, 1]^d)$ almost surely and the covariance operator of $X^{(d)}$ has the system of eigenvalues

$$(1.3) \quad \lambda_k^2 := \prod_{l=1}^d \lambda(k_l)^2, \quad k \in \mathbb{N}^d.$$

In the following we drop the index d and write $X(t)$ instead of $X^{(d)}(t)$. We set $T = [0, 1]^d$.

For any $n > 0$, let X_n be the partial sum of (1.2) corresponding to n maximal eigenvalues. We study the behavior of average error of approximation to X by X_n

$$e(X, X_n; d) = (\mathbb{E} \|X - X_n\|_{L_2(T)}^2)^{1/2},$$

as $d \rightarrow \infty$. Since in the following we consider only $L_2(T)$ -norms, we will simply write $\|\cdot\|$ instead of $\|\cdot\|_{L_2(T)}$. It is well known (see, for example, [2], [4] or [8]) that X_n provides the minimal average quadratic error among all linear approximations to X having rank n .

Since we are going to explore a *family* of random functions, it is more natural to investigate *relative* errors, that is to compare the error size with the size of the function itself.

Let

$$\Lambda := \sum_{i=1}^{\infty} \lambda(i)^2,$$

then

$$\mathbb{E} \|X\|^2 = \sum_{k \in \mathbb{N}^d} \lambda_k^2 = \Lambda^d.$$

Define the relative average approximation cardinality as

$$n(\varepsilon, d) := \min\{n : \frac{e(X, X_n; d)}{(\mathbb{E} \|X\|^2)^{1/2}} \leq \varepsilon\} = \min\{n : \mathbb{E} \|X - X_n\|^2 \leq \varepsilon^2 \Lambda^d\}.$$

The study of $n(\varepsilon, d)$ we are interested in here belongs to the class of problems dealing with the dependence of the complexity for linear multivariate problems on the dimension, see the works of H. Woźniakowski ([9], [10], [11], [12]) and the references therein.

It was suggested in [5] to use an auxiliary probabilistic construction for studying the properties of deterministic array of eigenvalues (1.3). We also follow this approach.

Consider a sequence of independent identically distributed random variables (U_l) , $l = 1, 2, \dots$ with the common distribution given by

$$(1.4) \quad \mathbb{P}(U_l = -\log \lambda(i)) = \frac{\lambda(i)^2}{\Lambda}, \quad i = 1, 2, \dots$$

Under assumption

$$(1.5) \quad \sum_{i=1}^{\infty} |\log \lambda(i)|^3 \lambda(i)^2 < \infty,$$

the condition $\mathbb{E}|U_l|^3 < \infty$ is obviously satisfied.

Let M and σ^2 denote the mean and the variance of U_l , respectively. Clearly,

$$\begin{aligned} M &= - \sum_{i=1}^{\infty} \log \lambda(i) \frac{\lambda(i)^2}{\Lambda}, \\ \sigma^2 &= \sum_{i=1}^{\infty} |\log \lambda(i)|^2 \frac{\lambda(i)^2}{\Lambda} - M^2. \end{aligned}$$

Then the third central moment of U_l is given by

$$\alpha^3 := \mathbb{E}(U_l - M)^3 = - \sum_{i=1}^{\infty} \log \lambda(i)^3 \frac{\lambda(i)^2}{\Lambda} - 3M\sigma^2 - M^3.$$

If (1.5) is verified, we have $|M| < \infty$, $0 \leq \sigma^2 < \infty$ and $|\alpha| < \infty$.

In the sequel the explosion coefficient

$$(1.6) \quad \mathcal{E} := \Lambda e^{2M}$$

will play a significant role. It was shown in [5] that $\mathcal{E} > 1$, except for the totally degenerate case when the number of strictly positive eigenvalues is zero or one. In other words, $\mathcal{E} = 1$ iff $\sigma = 0$. Henceforth we will exclude this degenerate case.

The following result was obtained in [5], Theorem 3.2.

Theorem 1.1 *Assume that the sequence $(\lambda(i))$ satisfies*

$$\sum_{i=1}^{\infty} |\log \lambda(i)|^2 \lambda(i)^2 < \infty.$$

Then for any $\varepsilon \in (0, 1)$ we have

$$\lim_{d \rightarrow \infty} \frac{\log n(\varepsilon, d) - d \log \mathcal{E}}{\sqrt{d}} = 2q,$$

where the quantile $q = q(\varepsilon)$ is chosen from the equation

$$(1.7) \quad 1 - \Phi\left(\frac{q}{\sigma}\right) = \varepsilon^2.$$

The authors of [5] conjectured that under further assumptions on the sequence $(\lambda(i))$ one can prove that

$$n(\varepsilon, d) \approx \frac{C(\varepsilon) \mathcal{E}^d e^{2q\sqrt{d}}}{\sqrt{d}}, \quad d \rightarrow \infty.$$

Indeed, we are going to confirm this conjecture.

2 Main result

It turns out that two different cases depending on the nature of the distribution of U_l should be distinguished. The proof and the final result rely on the fact whether this distribution is a *lattice* one or not.

Recall that one calls a discrete distribution of a random variable U a lattice distribution, if there exist numbers a and $h > 0$ such that every possible value of U can be represented in the form $a + \nu h$, where ν is an integer. The number h is called a span of the distribution. In the following, when studying the lattice case, we assume that h is a maximal span of the distribution, i.e. that it is impossible to represent all possible values of U_l in the form $b + \nu h_1$ for some b and $h_1 > h$.

Definition (1.4) yields that the variables U_l have a common lattice distribution iff $\lambda(i) = C e^{-n_i h}$ for some positive real C , h and $n_i \in \mathbb{N}$. We call this situation the *lattice case* and will assume that h is chosen to be the largest possible. Otherwise we say that the *non-lattice case* takes place.

Theorem 2.1 *Let the sequence $(\lambda(i))$, $i = 1, 2, \dots$ satisfy (1.5).*

Then for any $\varepsilon \in (0, 1)$ we have

$$n(\varepsilon, d) = K \phi\left(\frac{q}{\sigma}\right) \mathcal{E}^d e^{2q\sqrt{d}} d^{-1/2} (1 + o(1)), \quad d \rightarrow \infty,$$

where

$$\begin{aligned} \phi(x) &= \frac{1}{\sqrt{2\pi}} e^{-x^2/2}, \\ K &= \begin{cases} \frac{h}{\sigma(1-e^{-2h})} & \text{in the lattice case} \\ \frac{1}{2\sigma} & \text{otherwise,} \end{cases} \end{aligned}$$

and the quantile $q = q(\varepsilon)$ is defined in (1.7).

Remarks:

- One can see that the complexity of approximation increases exponentially as $d \rightarrow \infty$. This phenomenon is referred to as "dimensionality curse" or "intractability", see e.g. [8] and [10]. The notion of "curse of dimensionality" dates back at least to Bellman [1].

- Since by l'Hospital rule,

$$\lim_{h \rightarrow 0} \frac{h}{1 - e^{-2h}} = \frac{1}{2},$$

the formulae for K agree when $h \rightarrow 0$.

Proof:

Let $\zeta = \zeta(\varepsilon, d)$ be the maximal positive real number such that the sum of eigenvalues satisfies

$$\sum_{k \in \mathbb{N}^d: \lambda_k < \zeta} \lambda_k^2 \leq \varepsilon^2 \Lambda^d.$$

Define a lattice set in $\mathbb{N}^d$

$$A = A(\varepsilon, d) := \{k \in \mathbb{N}^d : \lambda_k \geq \zeta\} = \{k \in \mathbb{N}^d : \prod_{l=1}^d \lambda(k_l) \geq \zeta\}.$$

Since for any $k \in A$ it holds that $\lambda_k > 0$, one can write

$$\begin{aligned} n(\varepsilon, d) &= \text{card}(A) = \sum_{k \in A} \frac{\lambda_k^2}{\lambda_k^2} \\ &= \sum_{k \in \mathbb{N}^d: -\sum \log \lambda(k_l) \leq -\log \zeta} \Lambda^d \exp\{-2 \sum_{l=1}^d \log \lambda(k_l)\} \prod_{l=1}^d \mathbb{P}(U_l = -\log \lambda(k_l)) \\ &= \Lambda^d \mathbb{E} \exp\{2 \sum_{l=1}^d U_l\} \mathbb{I}_{\{\sum_{l=1}^d U_l \leq -\log \zeta\}}. \end{aligned}$$

For centered and normalized sums

$$Z_d = \frac{\sum_{l=1}^d U_l - dM}{\sigma \sqrt{d}}$$

we have

$$\left\{ \sum_{l=1}^d U_l \leq -\log \zeta \right\} = \{Z_d \leq \theta\},$$

where

$$(2.8) \quad \theta = \theta(\varepsilon, d) = -\frac{\log \zeta + dM}{\sigma \sqrt{d}}.$$

We show now that θ has a useful probabilistic meaning in terms of (U_l) and of their sums. Applying Lemma 3.1 from [5] we have for any $d \in \mathbb{N}$ and $z \in \mathbb{R}^1$

$$\begin{aligned} \sum_{k \in \mathbb{N}^d: \lambda_k < z} \lambda_k^2 &= \Lambda^d \mathbb{P} \left(\sum_{l=1}^d U_l > -\log z \right) \\ &= \Lambda^d \mathbb{P} \left(Z_d > -\frac{\log z + dM}{\sigma \sqrt{d}} \right) = \Lambda^d \mathbb{P}(Z_d > \theta_z), \end{aligned}$$

where

$$\theta_z = -\frac{\log z + dM}{\sigma\sqrt{d}}.$$

Fix $\varepsilon \in (0, 1)$. Observe that

$$\sum_{k \in \mathbb{N}^d: \lambda_k < z} \lambda_k^2 \leq \varepsilon^2 \Lambda^d$$

iff

$$\mathbb{P}(Z_d > \theta_z) \leq \varepsilon^2.$$

Therefore, $\theta = \theta(\varepsilon, d)$ defined by (2.8) is the $(1 - \varepsilon^2)$ -quantile of the distribution of Z_d , namely,

$$\theta(\varepsilon, d) = \min\{\theta : \mathbb{P}(Z_d > \theta) \leq \varepsilon^2\} = \min\{\theta : \mathbb{P}(Z_d \leq \theta) > 1 - \varepsilon^2\}.$$

Let $q = q(\varepsilon)$ be the quantile of the normal distribution function chosen from the equation (1.7). Then in view of the Central Limit Theorem

$$(2.9) \quad \theta(d, \varepsilon) \rightarrow \frac{q(\varepsilon)}{\sigma}, \quad d \rightarrow \infty,$$

for any fixed $\varepsilon \in (0, 1)$.

Now let us return to the average approximation cardinality. We obtain

$$\begin{aligned} n(\varepsilon, d) &= \mathcal{E}^d \mathbb{E} \exp\{2\sigma\sqrt{d}Z_d\} \mathbb{I}_{\{Z_d \leq \theta\}} \\ &= \mathcal{E}^d \exp\{2\sigma\sqrt{d}\theta\} \int_{-\infty}^{\theta} \exp\{2\sigma\sqrt{d}(z - \theta)\} dF_d(z), \end{aligned}$$

where $F_d(z) = \mathbb{P}(Z_d < z)$ and $\mathcal{E}$ is defined in (1.6).

Denote

$$\Psi_d(z) := \exp\{2\sigma\sqrt{d}(z - \theta)\}$$

and integrate by parts the integral

$$\int_{-\infty}^{\theta} \Psi_d(z) d[F_d(z) - F_d(\theta)] = \int_{-\infty}^{\theta} [-F_d(z) + F_d(\theta)] d\Psi_d(z).$$

Non-lattice case

In the following part of the proof we will assume that the distribution of (U_l) is not a lattice one. This is true in most interesting cases, such as the Brownian sheet, the fractional Brownian sheet, the Brownian pillow, the Brownian pillow-slip (Kiefer field).

In view of (1.5) we are able to apply Cramér-Esseen Theorem (cf. Theorem 2 §42 in [3], Theorem 5.21 §5.7 of Chapter V in [7] or Theorem 4 §3 of Chapter VI in [6]). It

leads to

$$\begin{aligned}
(2.10) \quad & \int_{-\infty}^{\theta} [-F_d(z) + F_d(\theta)] d\Psi_d(z) = \int_{-\infty}^{\theta} [-\Phi(z) + \Phi(\theta)] d\Psi_d(z) \\
& + \frac{\alpha^3}{6\sigma^3\sqrt{2\pi d}} \int_{-\infty}^{\theta} [(z^2 - 1)e^{-z^2/2} - ((\theta^2 - 1)e^{-\theta^2/2})] d\Psi_d(z) + o\left(\frac{1}{\sqrt{d}}\right) \\
& = I_1 + I_2 - I_3 - I_4 + o\left(\frac{1}{\sqrt{d}}\right),
\end{aligned}$$

where

$$\begin{aligned}
I_1 &= \int_{-\infty}^{\theta} [-\Phi(z) + \Phi(\theta)] d\Psi_d(z), \\
I_2 &= \frac{\alpha^3}{6\sigma^3\sqrt{2\pi d}} \int_{-\infty}^{\theta} z^2 e^{-z^2/2} d\Psi_d(z), \\
I_3 &= \frac{\alpha^3}{6\sigma^3\sqrt{2\pi d}} \int_{-\infty}^{\theta} e^{-z^2/2} d\Psi_d(z), \\
I_4 &= \frac{\alpha^3}{6\sigma^3\sqrt{2\pi d}} (\theta^2 - 1) e^{-\theta^2/2} \sim \frac{\alpha^3}{6\sigma^3\sqrt{2\pi d}} \left(\left(\frac{q}{\sigma} \right)^2 - 1 \right) e^{-q^2/2\sigma^2}, \quad d \rightarrow \infty.
\end{aligned}$$

The last equivalence is provided by (2.9).

Since $d\Psi_d(z) = 2\sigma\sqrt{d}\Psi_d(z)dz$, then I_2 is given, after a change of variable, by

$$I_2 = I_2(d, \theta) = \frac{\alpha^3}{3\sigma^2\sqrt{2\pi d}} \int_0^{\infty} \left(\theta - \frac{y}{\sqrt{d}} \right)^2 \exp\left\{ -\frac{1}{2} \left(\theta - \frac{y}{\sqrt{d}} \right)^2 \right\} \exp\{-2\sigma y\} dy$$

where $y = -\sqrt{d}(z - \theta)$.

For any $d = 1, 2, \dots$

$$0 \leq \left(\theta - \frac{y}{\sqrt{d}} \right)^2 \exp\left\{ -\frac{1}{2} \left(\theta - \frac{y}{\sqrt{d}} \right)^2 \right\} \leq (|\theta| + y)^2.$$

This estimate gives us the majorant required in Lebesgue Theorem. Using (2.9) and passing to the limit in the integral we obtain, as $d \rightarrow \infty$,

$$I_2(d, \theta) = \frac{\alpha^3}{6\sigma^3\sqrt{2\pi d}} \left(\frac{q}{\sigma} \right)^2 e^{-q^2/2\sigma^2} (1 + o(1)).$$

Similarly,

$$I_3(d, \theta) = \frac{\alpha^3}{6\sigma^3\sqrt{2\pi d}} e^{-q^2/2\sigma^2} (1 + o(1)).$$

Thus we obtain that $\sqrt{d}I_4 = \sqrt{d}(I_2 - I_3)(1 + o(1))$, hence, $I_2 - I_3 - I_4 = o\left(\frac{1}{\sqrt{d}}\right)$.

Consider the main integral I_1 .

$$\begin{aligned}
I_1 &= I_1(d, \theta) = \int_{-\infty}^{\theta} [-\Phi(z) + \Phi(\theta)] d\Psi_d(z) \\
&= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\theta} \exp\{2\sigma\sqrt{d}(z - \theta)\} \exp\{-z^2/2\} dz \\
&= \frac{1}{\sqrt{2\pi d}} \int_0^{\infty} \exp\{-\frac{1}{2}(\theta - \frac{y}{\sqrt{d}})^2\} \exp\{-2\sigma y\} dy \\
(2.11) \quad &\sim \frac{1}{2\sigma\sqrt{2\pi d}} e^{-q^2/2\sigma^2}, \quad d \rightarrow \infty.
\end{aligned}$$

Then

$$n(\varepsilon, d) = \frac{\mathcal{E}^d \exp\{2q\sqrt{d}\}}{2\sigma\sqrt{d}} \frac{1}{\sqrt{2\pi}} \exp\{-q^2/2\sigma^2\} (1 + o(1)),$$

as asserted.

Lattice case

Now we will proceed under assumption that the sequence (U_l) has a lattice distribution.

Let the possible values of the random variable U_l be

$$\tilde{a} + \nu h, \quad \nu = 0, \pm 1, \pm 2, \dots$$

where $\tilde{a} = M + a$ is the shift, and h is the maximal span of the distribution. Therefore, all possible values of Z_d have the form

$$\frac{da + \nu h}{\sigma\sqrt{d}}, \quad \nu = 0, \pm 1, \pm 2, \dots$$

Introduce the function

$$S(x) = [x] - x + \frac{1}{2},$$

where $[x]$ denotes, as usual, the integer part of x , and consider

$$S_d(x) = \frac{h}{\sigma} S\left(\frac{x\sigma\sqrt{d} - da}{h}\right).$$

Let $F_d(z)$ be as above. Then under assumption (1.5) Esseen's result (see Theorem 1 § 43 in [3]) yields

$$F_d(z) - \Phi(z) = \frac{e^{-z^2/2}}{\sqrt{2\pi}} \left(\frac{S_d(z)}{\sqrt{d}} - \frac{\alpha^3(z^2 - 1)}{6\sigma^3\sqrt{d}} \right) + o\left(\frac{1}{\sqrt{d}}\right)$$

uniformly in z .

Comparing with (2.10), we observe that one only needs to evaluate the additional term

$$\begin{aligned}
J &= \frac{1}{\sqrt{2\pi d}} \int_{-\infty}^{\theta} [-S_d(z)e^{-z^2/2} + S_d(\theta)e^{-\theta^2/2}] d\Psi_d(z) \\
&= \frac{1}{\sqrt{2\pi d}} \int_{-\infty}^{\theta} \Psi_d(z) d\left(S_d(z)e^{-z^2/2}\right) = J_1 - J_2 + J_3,
\end{aligned}$$

where

$$\begin{aligned} J_1 &= \frac{1}{\sqrt{2\pi d}} \int_{-\infty}^{\theta} \Psi_d(z) S'_d(z) e^{-z^2/2} dz, \\ J_2 &= \frac{1}{\sqrt{2\pi d}} \int_{-\infty}^{\theta} \Psi_d(z) S_d(z) z e^{-z^2/2} dz, \end{aligned}$$

and J_3 is a "discrete part", which is defined in the following way.

Notice that $S(x)$ is a periodic function with period one, therefore $S_d(x)$ possesses a period $h/\sigma\sqrt{d}$ and has jumps at points $\{\frac{kh+da}{\sigma\sqrt{d}}, k \in \mathbb{Z}\}$. If the point θ belongs to this lattice then there exists an integer k' such that $\theta = \frac{k'h+da}{\sigma\sqrt{d}}$. Hence, one can integrate the discontinuous part of the integral J with respect to the measure $\frac{h}{\sigma}\delta_{\frac{kh+da}{\sigma\sqrt{d}}}$ and obtain

$$J_3 = \frac{1}{\sqrt{2\pi d}} \frac{h}{\sigma} \sum_{k=-\infty}^{k'} \Psi_d\left(\frac{kh+da}{\sigma\sqrt{d}}\right) \exp\left\{-\frac{1}{2}\left(\frac{kh+da}{\sigma\sqrt{d}}\right)^2\right\}.$$

We start with the estimation of J_1 . At the points where the derivative $S'_d(z)$ makes sense, one can easily calculate that $S'_d(z) = \frac{h}{\sigma} S\left(\frac{z\sigma\sqrt{d}-da}{h}\right) = -\sqrt{d}$, therefore, similarly to the non-lattice case, by Lebesgue Theorem we have

$$\begin{aligned} J_1 &= \frac{-\sqrt{d}}{\sqrt{2\pi d}} \int_{-\infty}^{\theta} \exp\{2\sigma\sqrt{d}(z-\theta)\} \exp\{-z^2/2\} dz \\ &= \frac{-1}{\sqrt{2\pi d}} \int_0^{\infty} \exp\left\{-\frac{1}{2}\left(\theta - \frac{y}{\sqrt{d}}\right)^2\right\} \exp\{-2\sigma y\} dy \\ (2.12) \quad &\sim \frac{-1}{2\sigma\sqrt{2\pi d}} e^{-q^2/2\sigma^2}, \quad d \rightarrow \infty, \end{aligned}$$

and it yields $\sqrt{d}J_1 = -\sqrt{d}I_1(1 + o(1))$.

As for the integral J_2 , this one, as d is large enough, plays no role at all. Indeed,

$$\begin{aligned} J_2 &= \frac{1}{\sqrt{2\pi d}} \int_{-\infty}^{\theta} \exp\{2\sigma\sqrt{d}(z-\theta)\} S_d(z) z \exp\{-z^2/2\} dz \\ &= \frac{1}{\sqrt{2\pi d}} \frac{1}{\sqrt{d}} \int_0^{\infty} \exp\left\{-\frac{1}{2}\left(\theta - \frac{y}{\sqrt{d}}\right)^2\right\} \left(\theta - \frac{y}{\sqrt{d}}\right) S_d\left(\theta - \frac{y}{\sqrt{d}}\right) \exp\{-2\sigma y\} dy \\ &\leq \frac{3h}{2\sigma d\sqrt{2\pi}} \int_0^{\infty} \exp\left\{-\frac{1}{2}\left(\theta - \frac{y}{\sqrt{d}}\right)^2\right\} \left(\theta - \frac{y}{\sqrt{d}}\right) \exp\{-2\sigma y\} dy \\ &\sim \frac{3h}{4\sigma^2 d\sqrt{2\pi}} \left(\frac{q}{\sigma}\right)^2 e^{-q^2/2\sigma^2}, \quad d \rightarrow \infty. \end{aligned}$$

And, of course, $J_2 = o\left(\frac{1}{\sqrt{d}}\right)$.

Now we consider the most essential summand

$$\begin{aligned}
J_3 &= \frac{1}{\sqrt{2\pi d}} \frac{h}{\sigma} \sum_{k=-\infty}^{k'} \exp\{2\sigma\sqrt{d} \left(\frac{kh + da}{\sigma\sqrt{d}} - \theta \right)\} \exp\left\{-\frac{1}{2} \left(\frac{kh + da}{\sigma\sqrt{d}} \right)^2\right\} \\
&= \frac{1}{\sqrt{2\pi d}} \frac{h}{\sigma} \sum_{k=-\infty}^{k'} \exp\{2h(k - k')\} \exp\left\{-\frac{1}{2} \left(\frac{kh + da}{\sigma\sqrt{d}} \right)^2\right\} \\
&= \frac{1}{\sqrt{2\pi d}} \frac{h}{\sigma} \sum_{l=0}^{\infty} \exp\{-2hl\} \exp\left\{-\frac{1}{2} \left(\frac{(k' - l)h + da}{\sigma\sqrt{d}} \right)^2\right\} \\
&= \frac{1}{\sqrt{2\pi d}} \frac{h}{\sigma} \sum_{l=0}^{\infty} \exp\{-2hl\} \exp\left\{-\frac{1}{2} \left(\theta - \frac{lh}{\sigma\sqrt{d}} \right)^2\right\} \\
(2.13) \quad &\sim \frac{1}{\sigma\sqrt{d}} \frac{h}{(1 - e^{-2h})} \frac{1}{\sqrt{2\pi}} e^{-q^2/2\sigma^2}, \quad d \rightarrow \infty.
\end{aligned}$$

We obtain

$$\sqrt{d}J_3 = \sqrt{d} \frac{2h}{(1 - e^{-2h})} I_1 (1 + o(1)).$$

Putting together (2.11), (2.12) and (2.13), we obtain

$$n(\varepsilon, d) = \frac{\mathcal{E}^d e^{2q\sqrt{d}}}{\sigma\sqrt{d}} \frac{h}{(1 - e^{-2h})} \frac{1}{\sqrt{2\pi}} e^{-q^2/2\sigma^2} (1 + o(1)), \quad d \rightarrow \infty.$$

□

Acknowledgements

This paper was partially written when the author was visiting the Institut für Mathematische Stochastik, Georg-August-Universität, Göttingen. Special thanks are due to Professor M.A. Lifshits for formulation of the problem and constant encouragement and to Professor M. Denker for his support and for providing excellent working conditions. The author was supported by grants RFBR 05-01-00911 and RFBR-DFG 04-01-04000.

References

- [1] Bellman, R. *Adaptive Control Processes: a Guided Tour*. Princeton University, Princeton, 1961, 255p.
- [2] Buslaev, A. P., Seleznev, O. V. *On certain extremal problems in the theory of approximation of random processes*. East J. Approx., 1999, v. 5, no. 4, 467–481.
- [3] Gnedenko, B. V., Kolmogorov, A. N. *Limit Distributions for Sums of Independent Random Variables*. Addison-Wesley, Cambridge, 1954, ix+264 p (*in Russian*: GTTI, Moscow-Leningrad, 1949).

- [4] Kühn, Th., Linde, W. *Optimal series representation of fractional Brownian sheets*. Bernoulli, 2002, v. 8, no. 5, 669–696.
- [5] Lifshits, M. A., Tulyakova, E. V. *Curse of dimensionality in approximation of random fields*. Probab. Math. Stat., 2006, v. 26, no 1, 83–98.
- [6] (*in Russian*: Nauka, Moscow, 1972, 416 p.).
- [7] Petrov, V. V. *Limit Theorems of Probability Theory: Sequences of Independent Random Variables*. Oxford Studies in Probability, 4. Clarendon press, Oxford, 1995, ix+292 p. (*in Russian*: Nauka, Moscow, 1987, 320 p.).
- [8] Ritter, K. *Average-case Analysis of Numerical Problems*. Lecture Notes in Mathematics, 1733. Springer-Verlag, Berlin, 2000, x+254 p.
- [9] Woźniakowski, H. *Average case complexity of linear multivariate problems. Part 1: Theory. Part 2: Applications*. J. Complexity, 1992, v. 8, 337–372, 373–392.
- [10] Woźniakowski, H. *Tractability and strong tractability of linear multivariate problems*. J. Complexity, 1994, v. 10, 96–128.
- [11] Woźniakowski, H. *Tractability and strong tractability of multivariate tensor product problems*. J. of Computing and Information, 1994, v. 4, 1–19.
- [12] Woźniakowski, H. *Tractability of multivariate problems for weighted spaces of functions*. Approximation and Probability, Banach Center Publ., v. 72, 2006, 407–427.

Nora Serdyukova

St.Petersburg State University

198504 Stary Peterhof

Dept of Mathematics

and Mechanics

Bibliotechnaya pl., 2

Russia

`nora.serdyukova@ns14797.spb.edu`